

Fish Health Events (Quarter 4 2008)						
Fish Health SubZone	Species	Life Stage	Veterinary Diagnosis	Count of Fish Health Events ^{1,2,3}		
				New	Ongoing/Recurring	Relapsing
All	Atlantic Salmon	"Early"		0	0	0
All zones ⁵	Atlantic Salmon	"Later"		0	0	0
2-3	Atlantic Salmon	"Later"	Lepeophtheirus Infection	0	2	0
			Myxobacterial Infection	5	1	0
			Piscirickettsia salmonis Infection	2	2	0
2-4	Atlantic Salmon	"Later"	Lepeophtheirus Infection	2	0	0
			Myxobacterial Infection	1	0	0
3-1 + 3-2	Atlantic Salmon	"Later"	Lepeophtheirus Infection	3	0	0
			Myxobacterial Infection	2	1	0
3-3	Atlantic Salmon	"Later"	Aeromonas salmonicida (Atypical) Infection	1	0	0
			Lepeophtheirus Infection	9	0	0
			Viral Haemorrhagic Septicemia Virus Infection	1	0	0
			Myxobacterial Infection	0	1	0
3-4 + 3-5	Atlantic Salmon	"Later"	Lepeophtheirus Infection	0	2	0
All zones	Pacific Salmonids	"Early"	Aeromonas salmonicida Infection	1	0	0
			Case worked up but no diagnosis	0	1	0
All zones	Pacific Salmonids	"Later"	Renibacterium salmoninarum Infection	2	4	0
			Vibrio (Listonella) Infection	1	1	0

Notes

- 1 Reporting reflects life stage rather than water type. See notes 1 - 2 of Average Mortality Rate report.
- 2 Counts of veterinary diagnosis are based on FISH GROUP (not site); more than one fish group may exist at a site
- 3 Fish Health Events reflect the following categories:

New	First time occurrence; new event
Ongoing/recurring	Repeat or ongoing occurrence from previous calendar quarter
Relapsing	Repeat occurrence from calendar quarter at least two quarters preceding the current one
- 4 "Case worked up but no diagnosis" category requires workup and management steps taken, e.g., further investigation, husbandry change etc.
- 5 This field has been added to encompass a small number of later lifestage Atlantic salmon (e.g., broodstock) raised in areas other than the subzones shown above.